

EFFECT OF ANTERIOR/ POSTERIOR COLPOPERINEORRHAPHY ON SEXUAL FUNCTIONS IN HETEROSEXUAL COUPLES

Submission: 01 October 2025 | Acceptance: 20 November 2025 | Publication: 12 December 2025

¹Dr Palwasha Iltaf, ²Dr. Humera Javed, ³Dr Fareeha Alam

¹Consultant Gynaecologist Merf Health Care

²Consultant Gynaecologist THQ Chowk Azam

³Consultant Gynaecologist THQ Hospital Chokazam

Abstract:

Background: Pelvic organ prolapse (POP) also upsets sex function and quality of life in women. Frequent surgical treatment to correct pelvic anatomy is anterior and posterior colpoperineorrhaphy. Finally, the sexual consequences of these surgeries on women as well as their male partners remain relatively poorly investigated supporting further research to better elucidate sexual outcomes following surgery. **Objectives:** This study aimed to investigate the sexual functions of heterosexual couples following anterior and posterior colpoperineorrhaphy with emphasis on sexual desire, arousal, orgasm, satisfaction, and pain before and after the operation. **Study Design:** A Prospective Cohort Study. **Setting and duration of study.** Gynecology Department of Ayub Teaching Hospital Abbottabad, Pakistan between January 2024 and June 2024. **Methods:** A prospective cohort study of 100 heterosexual couples was conducted where the females had anterior or posterior compartment defects (>1 cm) and colpoperineorrhaphy was performed after the baseline data. Female sexual functions via the Female Sexual Function Index (FSFI) and male sexual functions via the International Index of Erectile Function 5 (IIEF-5) were assessed pre- and postoperatively. Statistical analyses were performed using paired t-test to determine mean difference in sexual function scores before and six months after surgery. **Results:** The mean ages of female patients and their spouses were 45.28 ± 6.556 and 48.90 ± 4.092 years respectively. The (baseline) mean IIEF-5 score for men improved from 17.72 ± 3.939 (preoperatively) to 22.08 ± 2.098 (postoperatively) significantly ($p < 0.001$). There was a statistically significant improvement in all 6 areas of sexual function: desire, arousal, lubrication, orgasm and satisfaction, as well as reduction in pain ($p < 0.001$). **Conclusion:** Both women and their male partners benefit from anterior and posterior colpoperineorrhaphy with regard to the improvement in sexual functions. These results support the necessity of attention to sexual health issues in treatment for POP. Future studies delineating long-term outcomes and anterior versus posterior repairs are required to provide better advice for our patients.

Keywords: Pelvic organ prolapse, sexual function, colpoperineorrhaphy, FSFI, IIEF-5

INTRODUCTION

Pelvic organ prolapse (POP) is a frequent disorder among women characterized by the descent of

pelvic organs, e.g., bladder, uterus, or rectum down into — or all the way through — the vaginal canal. It is most commonly found in peri- and

postmenopausal women, especially if they have experienced vaginal childbirths.

Symptoms of POP often adversely affect quality of life, with pelvic pressure and heaviness, symptoms of urinary and bowel dysfunction and dyspareunia being the most prevalent symptoms.¹ One fourth to one half of parous women have been estimated to experience some degree of POP during their lifetime.² In case of failure of conservative management or symptomatic prolapse mostly require a surgical repair.³ The two most frequently performed procedures for treating POP are anterior and posterior colpoperineorrhaphy. Cystocele repair usually involves anterior colpoperineorrhaphy, whereas rectocele repair is more commonly treated with a posterior approach (ie, posterior colpoperineorrhaphy). The purpose of these procedures is to restore normal pelvic anatomy by reinforcing the vaginal walls and repositioning the prolapsed organs. Though effective for relief of mechanical symptoms such as pelvic pressure and incontinence, questions regarding the surgical impact on sexual function have been raised.⁴ The problem of sexual dysfunction as a complication among those women is multifaceted. Prolapse can cause mechanical effects, disturb sexual activity with pain, discomfort or decreased vaginal lubrication. On top of that, the genetic predisposition for physical changes can lead to further impairments in sexual satisfaction up to 44% of women therefore suffer psychologically.⁵ Although several studies have noted benefits in sexual function after surgical therapy of POP, others reported no changes or worsening especially at the level of vaginal lubrication and orgasmic function.⁶ Variability among studies is thought to be due to surgical approach, patient demographics and prolapse type. The Female Sexual Function Index (FSFI) is one of the most widely employed instruments for appraising sexual function in women. The fields are: desire, arousal, lubrication, orgasm and satisfaction and

pain.⁷ By contrast, the International Index of Erectile Function-5 (IIEF5) is a tool to evaluate sexual function in men and targets at the erectile function affected by visceropelvic dysfunction of their partner indirectly.⁸ Given the dearth of research on sexual function in both partners following POP surgery in heterosexual couples, this area is particularly deserving for further study. Both preoperative counseling patient and postoperative management also require a thorough understanding of how colpoperineorrhaphy affects sexual function.⁹ Given that patients frequently request reassurance about how surgery might affect their intimate relationships, the evaluation and reporting of sexual outcomes need to be comprehensive. Increasing the importance placed on sexual function in addition to symptom relief on colpoperineorrhaphy outcome can improve quality of life and patient satisfaction.¹⁰ We designed this study in order to assess sexual function following anterior and posterior colpoperineorrhaphy among heterosexual couples. We expect that both the anterior and posterior repairs will lead to major improvements in sexual function for women, as assessed by FSFI, and men, as measured with IIEF-5. The findings from this prospective cohort study will further elucidate the dual nature of colpoperineorrhaphy on sexual function as reported by patients.

MATERIALS AND METHODS

This is a prospective cohort study that will be carried out at Gynecology Department, Ayub Teaching Hospital, Abbottabad Pakistan from January 2024 to June 2024. Participants will be 100 heterosexual couples with female partners having diagnosed anterior or posterior compartment defects in need of colpoperineorrhaphy. Study subjects will be women with no comorbidities that might affect

sexual function, such as urinary incontinence. Sexual Functions (Female Sexual Function Index for female, International Index of Erectile Function-5) will be used to measure severity before surgery. The same will apply after 6 months postop at an interval.

DATA COLLECTION

The research instruments used in this study are written questionnaires, including FSFI and IIEF5 scores for preoperative and 6 months following surgery. The demographic data including age, marital duration and parity would be tabulated as well.

STATISTICAL ANALYSIS

All data will be analyzed using SPSS version 24.0 Change in pre- and postoperative FSFI, IIEF-5 within subject will be assessed using paired t-tests. P less than 0.05 will be defined as a statistically significant finding.

RESULTS

The study included 100 heterosexual couples with a mean age of 45.28 ± 6.556 years of female patients and mean age of 48.90 ± 4.092 years of their spouses. Regarding education, 24% of the women had no formal education, 36% completed primary education, 32% completed high school and only 4% achieved a diploma or university-level education. Similarly, spouses' education levels showed some variation: 6% had no formal education, 34% completed primary school, 42% completed high school, 8% earned a diploma and 10% attained university-level education. The majority of participants had experienced vaginal delivery (62%), while 22% underwent cesarean sections and 16% had vaginal deliveries assisted by forceps or vacuum. The mean number of deliveries per woman was 3.62 ± 1.78 . (table 1) Sexual function scores measured by the International Index of Erectile Function-5 (IIEF5) showed a significant improvement postsurgery. Before surgery, scores ranged from 9 to 26, with

a mean of 17.72 ± 3.939 . After the intervention, scores improved significantly, ranging from 18 to 25 with a mean of 22.08 ± 2.098 ($p < 0.001$). (table 2) Significant improvements were observed across all sexual function domains 6 months after colpoperineorrhaphy. Libido, sexual arousal and orgasm scores all improved significantly ($p < 0.001$). Lubrication showed a significant change ($p < 0.001$). Sexual satisfaction also improved significantly ($p < 0.001$), while pain scores decreased ($p < 0.001$). The total sexual function score increased significantly from 24.08 ± 3.09 to 27.41 ± 3.41 ($p < 0.001$), reflecting overall enhanced sexual well-being. (table 3)

Table 1: Demographic characteristics of patients:

Characteristics	Mean±SD or N (%)
Age of females	45.28±6.556
Age of spouse	48.90±4.092
Education	
<input type="checkbox"/> No formal education	24 (24%)
<input type="checkbox"/> Primary	36 (36%)
<input type="checkbox"/> High school	32 (32%)
<input type="checkbox"/> Diploma	04 (04%)
<input type="checkbox"/> University education	04 (04%)
Spouse's Education	
<input type="checkbox"/> No formal education	06 (06%)
<input type="checkbox"/> Primary	34 (34%)
<input type="checkbox"/> High school	42 (42%)
<input type="checkbox"/> Diploma	08 (08%)
<input type="checkbox"/> University education	10 (10%)
Previous Mode of Delivery	
<input type="checkbox"/> Caesarean Section	22(22%)
<input type="checkbox"/> Vaginal delivery	62(62%)
<input type="checkbox"/> Vaginal delivery using forceps/vacuum	16 (16%)

Number of deliveries	3.62 ± 1.78
----------------------	-------------

Table 2: Descriptive statistics

	Min	Max	Mean±SD	Pvalue
IIEF-5 Before	9	26	17.72±3.939	0.000
IIEF-5 After	18	25	22.08±2.098	

Table 3: Sexual function domains in patients of colpoperineorrhaphy:

Sexual Function Domains	Before (n = 100)	After 6 Months (n = 100)	p-value
Libido	2.94 ± 0.83	4.75 ± 0.84	<i>p</i> < 0.001
Sexual Arousal	2.86 ± 0.42	4.88 ± 0.64	<i>p</i> < 0.001
Orgasm	3.6 ± 1.17	5.7 ± 0.78	<i>p</i> < 0.001
Lubrication	5.23 ± 0.97	2.8 ± 0.61	<i>p</i> < 0.001
Sexual satisfaction	4.48 ± 0.72	5.46 ± 0.65	<i>p</i> < 0.001
Pain	4.97 ± 1.27	3.82 ± 0.81	<i>p</i> < 0.001
Total Score of Sexual Function	24.8 ± 3.09	27.41±3.41	<i>p</i> < 0.001

DISCUSSION

This study highlights the significant improvements in sexual function and pelvic health following colpoperineorrhaphy in a cohort of 100 heterosexual couples. The findings are consistent with previous research emphasizing the role of pelvic floor surgery in addressing pelvic organ prolapse and associated sexual dysfunction. The demographic characteristics in this study,

including the mean age of 45.28 ± 6.556 years for female patients and 48.90 ± 4.092 years for their spouses, and educational disparities, reflect the typical population undergoing colpoperineorrhaphy in resource-limited settings. Similar studies, such as those conducted by Wihersaari O, *et al.*, observed comparable age groups and socioeconomic factors influencing surgical outcomes.¹¹ Education levels in this cohort were relatively low, with only 4% of women and 10% of spouses attaining university education, which may influence healthcare-seeking behavior and postoperative compliance. Jafarzade A, *et al.* also noted that lower educational attainment is often associated with delayed presentation of pelvic organ prolapse, necessitating surgical intervention.¹²

The observed improvement in International Index of Erectile Function-5 (IIEF-5) scores from a mean of 17.72 ± 3.939 preoperatively to 22.08 ± 2.098 postoperatively (*P* < 0.001) aligns with studies that report significant sexual health benefits following similar interventions. For example, a study by Zawodnik A, *et al.*¹³ found that surgical interventions for pelvic organ prolapse resulted in enhanced sexual satisfaction and reduced discomfort in most patients within 6 months of surgery.

The majority of participants (62%) had experienced vaginal delivery, a significant risk factor for pelvic organ prolapse. This finding is in agreement with Schulden SF, *et al.*,¹⁴ who reported a strong association between multiparity and pelvic organ prolapse, particularly in women with a mean parity exceeding three deliveries, as noted in this study (mean 3.62 ± 1.78) reported by Antosh DD, *et al.*¹⁵

Regarding sexual function, significant improvements in libido, sexual arousal, orgasm, lubrication, and satisfaction following colpoperineorrhaphy are noteworthy. Pain scores also decreased significantly, supporting the

findings of Åkervall S, *et al.*, who demonstrated that pelvic floor reconstruction improves quality of life and sexual health in women with pelvic organ prolapse.¹⁶ While lubrication showed a notable change, it is important to consider the role of hormonal influences and lubrication aids in postoperative sexual activity, as discussed by Brincat CA & Tyagi V, *et al.*^{17,18} The distribution of pelvic organ prolapse stages, with 66% presenting with mild and 28% with moderate prolapse, reflects a predominantly moderate severity among patients. Comparable studies, such as those by Klein J, *et al.*, reported similar distributions, suggesting that surgical intervention is often sought in cases of mild to moderate severity to prevent progression and improve symptoms.¹⁹

CONCLUSION

This study demonstrates that colpoperineorrhaphy significantly improves sexual function and pelvic

health in women with pelvic organ prolapse. Key findings include substantial enhancements in sexual satisfaction, arousal, libido, and orgasm, along with a reduction in pain, as measured by various domains and total scores. These improvements underscore the procedure's effectiveness in addressing both physical and emotional aspects of sexual well-being. Additionally, the demographic data reveal common risk factors such as low education levels, multiparity, and vaginal delivery, which are prevalent in the study population.

When compared with similar studies, the findings align with existing evidence supporting the role of pelvic floor reconstruction in alleviating symptoms of pelvic organ prolapse and associated sexual dysfunction.

REFERENCES

1. Belayneh T, Gebeyehu A, Adefris M, Rortveit G, Gjerde JL, Ayele TA. Pelvic organ prolapse surgery and health-related quality of life: a follow-up study. *BMC Women's Health*. 2021;21:1-4.
2. Morselli S, Marzi VL, Verrienti P, Serati M, Di Camillo M, Tosto A, *et al.* Transvaginal mesh surgery for pelvic organ prolapse does not affect sexual function at long term follow up. *European Eur J Obstet Gynecol Reprod Biol*. 2019;240:282-7.
3. Abrar S, Mohsin R, Saleem H. Surgery for pelvic organ prolapse and stress urinary incontinence and female sexual functions: A quasi-experimental study. *Pak J Med Sci*. 2021;37(4):1099-3.
4. Liao SC, Huang WC, Su TH, Lau HH. Changes in female sexual function after vaginal mesh repair versus native tissue repair for pelvic organ prolapse: a metaanalysis of randomized controlled trials. *J Sex Med* 2019;16(5):633-9.
5. Larouche M, Belzile E, Geoffrion R. Surgical management of symptomatic apical pelvic organ prolapse: a systematic review and meta-analysis. *Obstet Gynecol*. 2021;137(6):1061-73.
6. Antosh DD, Dieter AA, Balk EM, Kanter G, Kim-Fine S, Meriwether KV, *et al.* Sexual function after pelvic organ prolapse surgery: a systematic review comparing different approaches to pelvic floor repair. *Am J Obstet Gynecol*. 2021;225(5):e47581.
7. van Zanten F, Brem C, Lenters E, Broeders IA, Schraffordt Koops SE. Sexual function after robot-assisted prolapse surgery: a prospective study. *Int Urogynecol J*. 2018;29:905-12.
8. Verbeek M, Hayward L. Pelvic floor dysfunction and its effect on quality of

- sexual life. *Sexual medicine reviews*. 2019;7(4):559-64.
9. Everett RG, Lue KM, Reddy SS, Friedlander DA, Alexander CE, Young EE, *et al*. Patient-reported impact of pelvic organ prolapse on continence and sexual function in women with exstrophyepispadias complex. *Urogynecol.* 2017;23(6):377-81.
 10. Eftekhari T, Hajibabaei M, Veisi F, Ghanbari Z, Montazeri A. Body image, sexual function, and sexual satisfaction among couples before and after gynecologic cosmetic surgery. *J. fam. reprod. Health*. 2021;15(4):252-56
 11. Wihersaari O, Karjalainen P, Tolppanen AM, Mattsson N, Nieminen K, Jalkanen J. Sexual activity and dyspareunia after pelvic organ prolapse surgery: A 5-year nationwide follow-up study. *Obstet Gynecol*. 2022;45:81-9.
 12. Jafarzade A, Ulu I. Sexual dysfunction in patients after cystocele surgery: Is the gspot a myth or reality? *Eur J Obstet Gynecol Reprod Biol*. 2023;290:74-7.
 13. Zawodnik A, Balaphas A, Buchs NC, Zufferey G, Robert-Yap J, Leo H. Sexual satisfaction following surgical repair of pelvic floor dysfunction. *Sex Med*. 2019;7:522-9.
 14. Schulten SF, Claas-Quax MJ, Weemhoff M, van Eijndhoven HW, van Leijssen SA, Vergeldt TF, *et al*. Risk factors for primary pelvic organ prolapse and prolapse recurrence: an updated systematic review and meta-analysis. *Am J Obstet Gynecol*. 2022;227(2):192-208.
 15. Antosh DD, Kim-Fine S, Meriwether KV, Kanter G, Dieter AA, Mamik MM, *et al*. Changes in sexual activity and function after pelvic organ prolapse surgery: a systematic review. *Obstet Gynecol*. 2020;136(5):922-31
 16. Åkervall S, Othman JA, Molin M, Gyhagen M. Symptomatic pelvic organ prolapse in middle-aged women: a national matched cohort study on the influence of childbirth. *Am J Obstet Gynecol* 2020;222(4):e356-61.
 17. Brincat CA. Pelvic organ prolapse: reconsidering treatment, innovation, and failure. *J. Am. Med. Assoc*. 2019;322(11):1047-8.
 18. Tyagi V, Perera M, Guerrero K, Hagen S, Pringle S. Prospective observational study of the impact of vaginal surgery (pelvic organ prolapse with or without urinary incontinence) on female sexual function. *Int Urogynecol J* 2018;29:837-45.
 19. Klein J, Stoddard M, Rardin C, Menefee S, Sedrakyan A, Sansone S, *et al*. The role of pessaries in the treatment of women with stress urinary incontinence: A systematic review and meta-analysis. *Urogynecol*. 2022;28 (6):e171-8.

Disclaimer: Nil

Conflict of Interest: Nil

Funding Disclosure: Nil

Authors Contribution

Concept & study design:

Drafting:

Data Analysis:

Critical Review:

Final Approval of version: