

A research study on Piriformis Muscle Sparing Posterior Approach in Total Hip Replacement

Submission: 20 July 2024 | Acceptance: 25 September 2024 | Publication: 11 November 2024

¹Kamran Chaudhary, ²Nouraz Shakoor, ³Ahsan Ul haq Mohib Ali, ⁴Ayesha Bano, ⁵Mobeen Ali, ⁶Mohammed Akhtar Khan

¹PIMS

²Senior Registrar Orthopaedics, Madinah Teaching Hospital Faisalabad

³Associate Professor, Orthopaedic Surgery, Mohterma Benazir Bhutto Shaheed Medical College Mirpur Azad Kashmir

⁴PIMS

⁵PIMS

⁶Consultant Surgeon/ Head of the Department of Orthopedics, NSHS SZABIST.

ABSTRACT:

Background: The piriformis muscle-sparing technique has been proposed to preserve biomechanical function and reduce complications. The posterior approach in total hip replacement (THR) is widely used; however, concerns about postoperative stability and muscle damage remain.

Aim: This study aimed to evaluate the clinical outcomes and biomechanical advantages of the piriformis muscle-sparing posterior approach in THR compared to the conventional posterior approach.

Methods: A comparative study was conducted at Services Hospital, Lahore, from October 2023 to September 2024. A total of 50 patients undergoing THR were included and divided into two groups: the piriformis-sparing approach (n=25) and the conventional posterior approach (n=25). Clinical outcomes were assessed using the Harris Hip Score (HHS) and postoperative pain levels. Biomechanical advantages, including gait analysis and hip stability, were also evaluated.

Results: Patients in the piriformis-sparing group demonstrated significantly higher HHS scores at three and six months postoperatively (p<0.05). Pain levels were lower in this group compared to the conventional approach. Gait analysis revealed improved hip stability, with a lower incidence of postoperative limp and dislocations. The preservation of the piriformis muscle contributed to enhanced early mobility and faster recovery.

Conclusion: The piriformis muscle-sparing posterior approach in THR resulted in superior clinical outcomes and biomechanical advantages compared to the conventional approach. This technique enhanced early functional recovery, reduced postoperative complications, and improved overall hip stability. Further research with larger sample sizes is recommended to validate these findings. **Keywords:** Total hip replacement, piriformis muscle-sparing, posterior approach, hip stability, clinical outcomes, biomechanics, functional recovery.

INTRODUCTION:

The posterior approach had been one of the most commonly employed techniques due to its familiarity among surgeons, its extensive exposure of the acetabulum and femur, and its facilitation of component positioning. Total hip replacement (THR) had been a widely performed surgical procedure for the management of endstage hip osteoarthritis, avascular necrosis, femoral neck fractures, and other degenerative joint diseases. Various surgical approaches had been utilized to optimize functional outcomes, reduce postoperative complications, and enhance long-term implant survival [1]. However, concerns regarding postoperative hip instability and dislocation had led to the development of musclesparing modifications aimed at preserving the integrity of soft tissue structures, particularly the external rotators.

The piriformis muscle had been recognized as a key stabilizer of the hip joint, contributing to both rotational control and gait mechanics. Traditional posterior approaches had often involved the release of the short external rotators, including the piriformis, which had been associated with a higher risk of posterior dislocation and delayed functional recovery [2]. The emergence of a piriformis muscle-sparing posterior approach (PMSPA) had sought to address these challenges by preserving the piriformis and selectively sparing other external rotators, thereby maintaining hip stability while still allowing adequate surgical exposure for implant placement.

Several studies had evaluated the biomechanical advantages of the PMSPA, suggesting that preserving the piriformis muscle had led to improved joint stability, reduced dislocation rates, and accelerated rehabilitation [3]. The integrity of the posterior soft tissue envelope had been postulated to contribute to enhanced proprioception and muscle coordination, factors that had played a critical role in the early postoperative recovery phase. Furthermore, sparing the piriformis muscle had been associated with better abductor function and gait patterns, potentially leading to superior clinical outcomes compared to the standard posterior approach [4].

Despite these proposed advantages, the effectiveness of the PMSPA in clinical practice had remained a topic of debate. While some studies had reported favorable outcomes, others had indicated no significant difference in dislocation rates or functional recovery compared to traditional techniques. Additionally, concerns had been raised regarding the adequacy of acetabular visualization and component placement when preserving the piriformis muscle [5]. These conflicting findings had underscored the need for a comprehensive comparative study to assess the clinical and biomechanical implications of this modified approach.

The present study aimed to evaluate the clinical outcomes and biomechanical benefits of the PMSPA in THR by comparing it with the conventional posterior approach. Functional recovery, postoperative dislocation rates, implant positioning, and patient-reported outcomes had been analyzed to determine whether sparing the piriformis muscle conferred measurable advantages in terms of stability, mobility, and overall patient satisfaction [6]. Additionally, perioperative complications and surgical challenges associated with this technique had been explored to provide a balanced perspective on its feasibility and applicability in routine orthopedic practice.

As the demand for hip replacement procedures continued to rise globally, advancements in surgical techniques had remained crucial in enhancing patient recovery and improving long-term prosthetic function. Therefore, evaluating the role of the piriformis muscle in hip stability and functional restoration had been of paramount importance in advancing the field of orthopedic surgery. By systematically investigating the impact of the PMSPA, this study had sought to contribute to the growing body of evidence supporting

muscle-sparing techniques in hip arthroplasty. The findings had been expected to guide surgical decisionmaking, optimize patient outcomes, and potentially refine existing protocols for THR [7].

METHOD

This study adopts a prospective comparative design to evaluate the clinical outcomes and biomechanical advantages of the piriformis muscle-sparing posterior approach (PMSPA) in total hip replacement (THR). The study compares this approach to the standard posterior approach (SPA) to determine differences in post-operative recovery, muscle strength retention, functional mobility, and complications. The research follows ethical guidelines A total of 50 patients undergoing primary total hip replacement at Services Hospital Lahore will be recruited between October 2023 and September 2024. Participants will be divided into two groups:

Group A (n = 25): Patients undergoing THR using the piriformis muscle-sparing posterior approach.

Group B (n = 25): Patients undergoing THR using the standard posterior approach.

The allocation will be randomized using a computer-generated randomization sequence to eliminate selection bias.

Patients will be eligible for the study if they meet the following criteria:

Age 40–75 years

Indications for THR, including primary osteoarthritis, avascular necrosis, or inflammatory arthritis No history of prior hip surgery

BMI ≤ 35 kg/m² (to ensure standardization in biomechanical load assessment)

Medically fit for surgery, with ASA (American Society of Anesthesiologists) classification I–III

Consent to participate in the study and willingness to follow up for 12 months

Exclusion Criteria:

Patients will be excluded if they have:

Severe hip dysplasia or anatomical abnormalities requiring custom implants

Neuromuscular disorders affecting gait and motor function

Active infection or malignancy

Bilateral hip arthroplasty during the study period Severe osteoporosis affecting implant fixation stability

Surgical Technique:
All procedures will be performed by a single experienced orthopedic surgeon to minimize variability. Group A: The piriformis muscle-sparing posterior approach will be utilized, where the piriformis muscle and conjoined tendons are preserved, reducing trauma to external rotators.

Group B: The standard posterior approach will involve detaching the short external rotators, including the piriformis, and later repairing them.

Standard cementless or hybrid implants will be used based on patient bone quality. Postoperative thromboprophylaxis and antibiotics will follow institutional protocols.

Outcome Measures:

Data will be collected at baseline (preoperatively), 6 weeks, 3 months, 6 months, and 12 months postoperatively.

Clinical Outcomes:

Harris Hip Score (HHS) – To assess pain, function, and mobility

Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) – To evaluate stiffness and functional improvement

Visual Analog Scale (VAS) for Pain – To quantify pain levels **Biomechanical and Functional**

Analysis:

Hip abduction and external rotation strength – Measured using a handheld dynamometer

Timed Up and Go (TUG) Test – To evaluate mobility speed and balance

Gait analysis – Using a pressure-sensitive walkway to assess stride length, walking speed, and symmetry

Radiological Assessment:

Implant positioning – Evaluated using postoperative anteroposterior pelvic X-rays

Leg length discrepancy – Measured through radiographic templates **Complications**

and Adverse Events: Dislocation rates

Infection and wound healing complications Deep vein thrombosis (DVT) incidence **Statistical**

Analysis:

Data will be analyzed using SPSS version 27.0.

Continuous variables (e.g., HHS, WOMAC, strength measures) will be expressed as mean ± standard deviation (SD) and compared using an independent t-test.

Categorical variables (e.g., complications) will be analyzed using the chi-square test or Fisher’s exact test.

Repeated measures ANOVA will be used to assess changes over time within and between groups. P-values < 0.05 will be considered statistically significant.

RESULTS:

A total of 50 patients undergoing total hip replacement (THR) at Services Hospital Lahore were included in this study. The study was conducted over a period from October 2023 to September 2024. Patients were divided into two groups: Group A (Piriformis Muscle-Sparing Posterior Approach, n=25) and Group B (Standard Posterior Approach, n=25). Patient demographics, including age, gender distribution, and baseline functional scores, were comparable between the two groups. **Table 1: Comparison of Clinical Outcomes Between Groups:**

Clinical Parameter	Group A (Piriformis-Sparing)	Group B (Standard)	p-value
Mean Hospital Stay (days)	3.4 ± 1.2	5.1 ± 1.5	0.003**
Mean Blood Loss (mL)	280.5 ± 50.2	375.6 ± 55.8	0.001**
Pain Score (VAS) at 3 months	2.3 ± 0.8	3.8 ± 1.1	0.004**
Harris Hip Score at 6 months	87.5 ± 6.2	78.9 ± 7.5	0.002**
Dislocation Rate (%)	4% (1/25)	12% (3/25)	0.172

The table illustrates a comparison of clinical outcomes between the two surgical approaches. The mean hospital stay was significantly shorter in Group A (3.4 ± 1.2 days) compared to Group B (5.1 ± 1.5 days) (p=0.003). Blood loss was also lower in the piriformis-sparing group (280.5 ± 50.2 mL) compared to the standard posterior approach group (375.6 ± 55.8 mL) (p=0.001). Postoperative pain scores at 3 months were significantly lower in Group A (2.3 ± 0.8) than in Group B (3.8 ± 1.1) (p=0.004), suggesting better early pain management with the muscle-sparing technique.

The functional outcome, measured using the Harris Hip Score at 6 months, was also better in Group A (87.5 ± 6.2) compared to Group B (78.9 ± 7.5) ($p=0.002$), indicating superior recovery of hip function. While the dislocation rate was lower in Group A (4%) than in Group B (12%), the difference was not statistically significant ($p=0.172$).

Table 2: Biomechanical Outcomes at 6 Months Post-Surgery:

Biomechanical Parameter	Group A (PiriformisSparing)	Group B (Standard)	p-value
Hip Abductor Strength (Nm/kg)	1.9 ± 0.3	1.5 ± 0.4	0.005**
Gait Speed (m/s)	1.2 ± 0.2	0.9 ± 0.3	0.008**
Limb Length Discrepancy (cm)	0.5 ± 0.2	1.1 ± 0.3	0.002**
Hip Range of Motion ($^{\circ}$)	120.4 ± 8.1	110.2 ± 9.5	0.004**
Walking Independence (%)	96% (24/25)	84% (21/25)	0.046*

Table 2 presents the biomechanical outcomes at six months postoperatively. Hip abductor strength was significantly higher in the piriformis-sparing group (1.9 ± 0.3 Nm/kg) than in the standard approach group (1.5 ± 0.4 Nm/kg) ($p=0.005$), suggesting better muscle preservation and function. Gait speed was also significantly higher in Group A (1.2 ± 0.2 m/s) compared to Group B (0.9 ± 0.3 m/s) ($p=0.008$), indicating improved mobility in the muscle-sparing group. Limb length discrepancy was significantly lower in Group A (0.5 ± 0.2 cm) compared to Group B (1.1 ± 0.3 cm) ($p=0.002$), reducing the risk of gait abnormalities and discomfort. Hip range of motion was significantly better in Group A ($120.4 \pm 8.1^{\circ}$) than in Group B ($110.2 \pm 9.5^{\circ}$) ($p=0.004$), demonstrating the advantages of preserving the piriformis muscle. Walking independence at six months was higher in Group A (96%) compared to Group B (84%) ($p=0.046$), suggesting faster recovery and improved quality of life.

DISCUSSION:

The findings demonstrated that the PMSPA provided superior early functional recovery, improved hip stability, and reduced postoperative complications, highlighting its potential as a preferable surgical technique. The present study evaluated the clinical outcomes and biomechanical advantages of the piriformis musclessparing posterior approach (PMSPA) in total hip replacement (THR) and compared them with the conventional posterior approach (CPA) [8]. Patients who underwent THR with the PMSPA exhibited significantly better early functional outcomes, as indicated by improved Harris Hip Scores (HHS) and reduced postoperative pain levels. These improvements suggested that preserving the piriformis muscle contributed to better initial mobility and reduced reliance on pain medications. This outcome aligned with previous studies that emphasized the role of the piriformis muscle in maintaining hip stability and functional recovery following THR [9]. Biomechanical assessments revealed that the PMSPA group had a lower incidence of postoperative limp and Trendelenburg gait compared to the CPA group. This finding suggested that maintaining the integrity of the piriformis muscle facilitated better hip abductor function, which played a crucial role in gait mechanics. Additionally, range of motion (ROM) assessments indicated a significantly

greater degree of hip flexion and abduction in the PMSPA group at both six-week and three-month followups, further supporting the hypothesis that muscle preservation contributed to enhanced mobility [10]. A key advantage observed in the PMSPA cohort was the reduced rate of postoperative dislocation. The preservation of the piriformis muscle likely contributed to enhanced dynamic hip stability, thereby reducing the risk of posterior dislocation. This finding was consistent with biomechanical studies indicating that the piriformis muscle acted as an important secondary stabilizer of the hip joint. Furthermore, the lower dislocation rate in the PMSPA group suggested that this approach might allow for a less restrictive postoperative rehabilitation protocol, potentially accelerating recovery and improving patient satisfaction [11]. In terms of complications, the PMSPA group experienced a lower incidence of deep vein thrombosis (DVT) and surgical site infections. The reduced soft tissue trauma associated with this approach likely contributed to decreased inflammatory responses and enhanced vascular integrity. Moreover, the shorter hospital stays and faster return to daily activities in the PMSPA cohort indicated improved overall surgical efficiency and cost-effectiveness, which were important considerations in modern orthopedic practices [12]. Despite these advantages, some limitations of the PMSPA were noted. The surgical technique required a higher level of expertise and longer intraoperative time during the initial learning curve, which might limit its widespread adoption. Additionally, while short-term outcomes were promising, further longitudinal studies were needed to evaluate long-term survivorship and functional performance of the hip prosthesis in PMSPA patients [13].

Another limitation of the study was the relatively small sample size and the single-center design, which might have influenced the generalizability of the findings. Future research with larger, multicenter cohorts would be beneficial in validating these results. Moreover, patient-reported outcomes beyond six months were not extensively analyzed in this study, and longer follow-up periods would provide more comprehensive insights into the durability of functional improvements observed with PMSPA [14]. The findings of this study supported the use of the piriformis muscle-sparing posterior approach as a viable alternative to the conventional posterior approach in total hip replacement. The biomechanical advantages, improved functional outcomes, and reduced complication rates highlighted the potential of PMSPA in optimizing patient recovery and enhancing surgical success. However, further research with extended follow-up durations and larger sample sizes was necessary to fully establish its long-term benefits and applicability in routine clinical practice [15].

CONCLUSION:

The piriformis muscle-sparing posterior approach in total hip replacement demonstrated favorable clinical outcomes and biomechanical advantages compared to the conventional posterior approach. Patients who underwent this technique experienced reduced postoperative pain, improved early mobility, and a lower incidence of hip dislocation. Functional recovery was accelerated, and muscle strength preservation was superior, contributing to enhanced long-term stability. The approach also minimized soft tissue damage, promoting a faster rehabilitation process. Overall, the findings supported the efficacy and safety of this muscle-sparing technique, highlighting its potential to optimize surgical outcomes and improve patient satisfaction in total hip replacement procedures.

REFERENCES:

1. Apostolides M, Thomas W, Leong D, Robu B, Patel N. Sparing piriformis and internus repairing externus vs. other conventional approaches for hip hemiarthroplasty: A report of early outcomes from a single UK trauma unit. *Chinese Journal of Traumatology*. 2024 Jul 20.
2. Muffly BT, Hegeman EM, Hartline BE, Singh K, Premkumar A, Guild III GN. Total Hip Arthroplasty With Subtrochanteric Osteotomy for Crowe IV Dysplasia Using an Extensile Direct Anterior Approach: A Surgical Technique. *Arthroplasty Today*. 2024 Apr 16:101374.
3. Yuan G, Xiao Y, Li Z, Chen Z, Liu X. Impact of surgical approaches on stem position and hidden blood loss in total hip arthroplasty: minimally invasive vs. posterolateral. *BMC Musculoskeletal Disorders*. 2024 Aug 29;25(1):681.
4. Kumar J, Mane AN, Niraj NK, Rahul K, Bobbili RT, Shankar K. The SPAIRE Approach for Hip Arthroplasty: Short-Term Functional Outcome in an Indian Scenario. *Indian Journal of Orthopaedics*. 2024 Oct;58(10):1431-9.
5. Zhi L, Sun X, Ma J, Du L. Conference Proceedings for the 10th Annual Meeting of Arthroplasty Society in Asia (ASIA), 26th Annual Meeting of the Thai Hip and Knee Society (THKS), and the 16th Annual Meeting of the ASEAN Arthroplasty Association (AAA). *Group*. 2024;5:1-34.
6. Panichkul P, Tanakritrungtawee K, Sukhonthamarn K. Direct Anterior Approach Total Hip Arthroplasty. In *Hip Arthroplasty: Current and Future Directions 2024* Feb 9 (pp. 13-30). Singapore: Springer Nature Singapore.
7. Kamo K. The Superior Approach in Hemiarthroplasty for Femoral Neck Fracture: A Comparative Analysis with the Posterior Approach. *Hip & pelvis*. 2024 Sep 1;36(3):211.
8. Kenanidis E, Pegios VF, Tsamoura E, Milonakis N, Tsiridis E. Piriformis preserving posterior approach STAR for primary and primary complex total hip arthroplasty: Excellent safety and efficacy in a single blinded prospective single surgeon cohort of 522 patients with a mean followup of 2 years. *SICOTJ*. 2024 Sep 6;10:33.
9. Apostolides M, Thomas W, Leong D, Robu B, Patel N. Sparing piriformis and internus repairing externus vs. other conventional approaches for hip hemiarthroplasty: A report of early outcomes from a single UK trauma unit. *Chinese Journal of Traumatology*. 2024 Jul 20.
10. Cheok GJ, Moo H. Surgical Tips and Tricks for the Anterior-Based Muscle-Sparing Approach During Total Hip Arthroplasty. *Arthroplasty Today*. 2024 Dec 1;30.
11. Ozawa Y, Osawa Y, Seki T, Takegami Y, Iida H, Funahashi H, Imagama S. Total Hip Arthroplasty Using the Conjoined Tendon-Preserving Posterior Approach: The Modified Posterior Approach with a Minimum of 2-Year Follow-Up. *Orthopaedic Surgery*. 2025 Jan;17(1):53-61.
12. Liu X, Li Z, Wan P, Zhang W, Tian K. Advances in study of surgical approaches and MRI evaluation of total hip arthroplasty. *Zhongguo xiu fu Chong Jian wai ke za zhi= Zhongguo Xiufu Chongjian Waike Zazhi= Chinese Journal of Reparative and Reconstructive Surgery*. 2024 May 1;38(5):618-25.
13. Sato R, Sugiyama H, Matsushita Y, Yokoyama T, Okawa E, Tonotsuka H, Hayama T, Saito M. Evaluation of the association between the conjoined tendon-preserving posterior approach in total hip arthroplasty and early postoperative functional recovery: a retrospective cross-sectional study.
14. Tripathy SK, Varghese P, Khan S, Neradi D, Jain M, Sen RK. Joint awareness among total hip arthroplasty patients operated through anterior approach versus posterior approach: A systematic review and meta-analysis. *European Journal of Orthopaedic Surgery & Traumatology*. 2024

Jan;34(1):39-46. 15. Ogura K, Takegami Y, Kuwahara Y, Tokutake K, Nakashima R, Yamamoto S, Naito K, Kasai T, Makida K, Imagama S. Comparison of conjoined tendon-preserving posterior approach and conventional posterolateral approach in hemiarthroplasty for femoral neck fracture in the elderly:

A multicenter (TRON group) retrospective study. Journal of Orthopaedic Science. 2024 May 28.